

# A Close-to-Aromatize Approach for the Late-Stage Functionalization through Ring Closing Metathesis

Boris Lozhkin<sup>a</sup> and Thomas R. Ward<sup>\*a</sup>

<sup>a</sup> Department of Chemistry, University of Basel, Building 1096, Mattenstrasse 24a, Biopark Rosental, CH-4058 Basel, Switzerland, e-mail: Thomas.ward@unibas.ch

© 2021 The Authors. Helvetica Chimica Acta published by Wiley-VHCA AG. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

An efficient approach for the synthesis of monosubstituted aromatic compounds relying on a ring-closing metathesis followed by spontaneous 1,2-elimination is presented. The efficiency for late-stage functionalization is highlighted in various solvents (up to 920 TON). This approach is compatible with strained cycles and other multiple bonds in the substrate.

**Keywords:** elimination, late-stage functionalization, metathesis, ring closing metathesis.

## Introduction

Aromatic fragments are present in a vast majority of organic compounds. For the late-stage introduction of aromatic fragments, common methods include various cross-coupling and C–H activation strategies.<sup>[1–6]</sup> Herein, we exploit ring closing metathesis of a readily accessible triene-ol to access monosubstituted benzene derivatives.


Numerous strategies to produce aromatic rings as a result of ring-closing metathesis (RCM) have been reported.<sup>[7–14]</sup> Most of these, however, require harsh oxidants to produce the aromatic ring.<sup>[15–20]</sup> We were inspired by a report by *Sabatino et al.*<sup>[21]</sup> who reported a ‘close-to-release’ approach whereby aromatization resulting from a 1,4-elimination serves as driving force for the generation of naphthalene, with the concomitant release of a cargo-molecule. Herein we report our efforts to extend this methodology to RCM followed by the 1,2-elimination of water to generate monosubstituted benzene moieties (*Scheme 1*).

## Results and Discussion

To test the generality of this approach, we identified the triene-one **1** as a key building block. Subsequently, a *Grignard*-type addition to the ketone functionality allows tethering this masked benzene to a variety of fragments.


Synthesis of (5*Z*,7*E*)-nona-1,5,7-trien-4-one **1** was carried out according to the general procedure reported by *Cossy* and coworkers that they used in the total synthesis of (–)-*muricatacin* and related compounds.<sup>[22]</sup> Reaction of *Weinreb* amide **2** with allyl-magnesium chloride afford triene **1** in >75% yield (*Scheme 2*). The  $\gamma,\delta$ -double bond is highly reactive, which, together with the (*Z*)-configuration of the  $\alpha,\beta$ -double bond, opens up the possibility of an intramolecular reaction.

The biphenyl precursor **3a** was selected for the optimization of the cyclization conditions. The reaction proceeds in low yields in polar solvents (*Table 1*, *Entries 1–3*), including in water using AquaMet as catalyst (*Table 1*, *Entries 4–6*). This may be traced back to



**Scheme 1.** Concomitant ring-closing metathesis followed by 1,2-elimination of water for the synthesis of monosubstituted benzene derivatives.

Supporting information for this article is available on the WWW under <https://doi.org/10.1002/hlca.202100024>



**Scheme 2.** Synthesis of (5Z,7E)-nona-1,5,7-trien-4-one **1**.


the very low solubility of the triene **3a** in aqueous media. The most effective solvent proved to be dichloromethane, in which all reagents are readily soluble. The

reaction yield is affected by the nature of the catalyst, with the *Hoveyda-Grubbs* second-generation (HG-II) performing best (Table 1, Entries 7–14). Strikingly, the yield of biphenyl **3** increases with decreasing catalyst loading from 5 mol-% to 1 mol-%. This can be traced back to the cross-metathesis between the diolefinic substrate **3a** and the *ortho*-isopropoxy-benzylidene moiety of HG-II catalyst. The yield of this cross-metathesis product is affected by catalyst loading. The resulting disubstituted olefin does not further participate in productive ring-closing metathesis, thus contributing to erode the yield of biphenyl **3** with increasing catalyst loading.

Near quantitative conversion was achieved using 0.1% catalyst, corresponding to >900 TON. <sup>1</sup>H-NMR Monitoring reveals that the equilibrium is reached after six hours at room temperature using 5 mol-% HG-II (Figure 1). To reach (near) full conversion, nine and twenty hours of reaction time are necessary using 1 mol-% and 0.1 mol-% HG-II, respectively (Figure 1).


With these optimized conditions at hand, monosubstituted benzenes **3–10** were synthesized without the requirement of isolating the alcohol resulting from the *Grignard* addition to (5Z,7E)-nona-1,5,7-trien-4-one (**1**). The syntheses were carried out in two steps: 1) the reaction of ketone **1** with a *Grignard* derivative and

**Table 1.** Optimization of cyclization conditions.




Entry <sup>[a]</sup>	Solvent	Catalyst	Cat. load. [%]	yield [%] <sup>[b]</sup>	TON
1	DMSO	HG-II	5	5 ± 1	1 ± 0
2	ACN	HG-II	5	20 ± 4	4 ± 1
3	Acetone	HG-II	5	26 ± 6	5 ± 2
4	DMSO/H <sub>2</sub> O	AquaMet	5	n.d.	0
5	ACN/H <sub>2</sub> O	AquaMet	5	3 ± 0	1 ± 0
6	Acetone/H <sub>2</sub> O	AquaMet	5	9 ± 4	2 ± 1
7	CH <sub>2</sub> Cl <sub>2</sub>	G-I	5	58 ± 5	12 ± 1
8	CH <sub>2</sub> Cl <sub>2</sub>	G-II	5	63 ± 6	13 ± 1
9	CH <sub>2</sub> Cl <sub>2</sub>	HG-I	5	71 ± 9	14 ± 2
10	CH <sub>2</sub> Cl <sub>2</sub>	Umicore-M1	5	57 ± 7	11 ± 1
11	CH <sub>2</sub> Cl <sub>2</sub>	Schrock <sup>[c]</sup>	5	35 ± 1	7 ± 0
12	CH <sub>2</sub> Cl <sub>2</sub>	HG-II	5	92 ± 3	18 ± 1
13	CH <sub>2</sub> Cl <sub>2</sub>	HG-II	1	97 ± 2	97 ± 2
14	CH <sub>2</sub> Cl <sub>2</sub>	HG-II	0.1	92 ± 4	920 ± 40

<sup>[a]</sup> Reaction conditions: solvent (1 M), catalyst, r.t., 6 h. <sup>[b]</sup> Yields were determined by <sup>1</sup>H-NMR as the average of three independent experiments. <sup>[c]</sup> 2,6-Diisopropylphenylimidoneophylidene molybdenum(VI) bis(hexafluoro-*tert*-butoxide).



**Figure 1.**  $^1\text{H}$ -NMR Monitoring of the cyclization ( $\text{CH}_2\text{Cl}_2$ , 1 M), 5 mol-% HG-II, r.t.). None of the diene-ol intermediates is detected by  $^1\text{H}$ -NMR.



**Scheme 3.** Evaluating the regioselectivity of RCM using tetraene-ol **11a**.

2) ring-closing metathesis with spontaneous 1,2-elimination of  $\text{H}_2\text{O}$  (Table 2).


The metallated substrates were obtained from the corresponding halides and alkynes by adding either Mg or BuLi in ether. The monosubstituted benzene products **3–10** resulting from RCM and 1,2-elimination were isolated by column chromatography. Gratifyingly, the above reaction sequence tolerates a variety of (cyclic) aliphatic and aliphatic substituents, including cyclobutane (Table 2, Entries 1–6). Alkynes do not participate in metathesis as no product mixtures could be detected (Entries 7 and 8).

The regioselectivity was evaluated by subjecting the tetraene-ol substrate **11a** to RCM. Under standard conditions ( $\text{CH}_2\text{Cl}_2$ , 1 M), 5 mol-% HG-II, r.t.), no RCM product was detected. Upon increasing catalyst loading to 50%, only cross-metathesis with the styrene fragment of the HG-II catalyst could be isolated, along with the starting material **11a**. We could not find any rationale for this unexpected lack of reactivity (Scheme 3).

## Conclusions

An efficient method for the synthesis of monosubstituted aromatic compounds with saturated and unsaturated

**Table 2.** Substrate scope for the synthesis of monosubstituted benzene derivatives through RCM.



Entry <sup>[a]</sup>	RMX	Product	Yield [%] (over 2 steps) <sup>[b]</sup>
1	MeMgI	<b>4</b>	65
2	BuMgBr	<b>5</b>	78
3	cyclobutyl- $\text{CH}_2\text{MgBr}$	<b>6</b>	56
4	BnMgBr	<b>7</b>	75
5	PhMgBr	<b>3</b>	83
6	<i>p</i> -MeOPhMgBr	<b>8</b>	72
7	$\text{HC}\equiv\text{CCH}_2\text{MgBr}$	<b>9</b>	70
8	$\text{PhC}\equiv\text{CLi}$	<b>10</b>	47

<sup>[a]</sup> The first step was carried out with ketone **1** and corresponding Grignard reagent (diethyl ether (0.5 M),  $0^\circ\text{C}$ , 2 h) without isolating the corresponding trien-4-ol. The second step was performed with crude alcohols using HG-II catalyst (5 mol-%) in  $\text{CH}_2\text{Cl}_2$  (1 M) for 6 h. <sup>[b]</sup> Yield of isolated product.

aliphatic and aromatic substituents was developed. The key step proceeds in almost quantitative yields with low catalyst loading (0.1 mol-%, up to 920 TON). This method can be used to introduce an aromatic moiety at a late stage if alternative cross-coupling schemes prove challenging. The synthesis of the key intermediate can be carried out on a multi-gram scale with a total yield of >52% over five steps. The strategy outlined herein, relying on more elaborate Weinreb amides,<sup>[22,23]</sup> should allow to introduce various poly-substituted benzene moieties as a result of ring-closing metathesis. The strategy to introduce a masked benzene moiety as a result of RCM is limited by the following features: 1) The substrates must not contain any functional group that can interfere with the metalating reagent and 2) The

metalated derivative must have sufficient nucleophilicity to react with the carbonyl group of the unsaturated ketone.

We are currently capitalizing on this strategy to generate drugs *in vivo* (e.g. tamoxifen) as a result of ring-closing metathesis from inactive precursors.<sup>[21]</sup>

## Acknowledgements

This project has received funding from the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie (grant agreement no. 765497 (THERACAT)).

## Author Contribution Statement

T. R. W. and B. L. designed the concept. T. R. W. supervised the work which was carried out by B. L.; B. L. and T. R. W. discussed the results and wrote the manuscript. Both authors have given approval to the final version of the manuscript.

## References

- [1] Y. Zhu, M. Bauer, J. Ploog, L. Ackermann, 'Late-Stage Diversification of Peptides by Metal-Free C–H Arylation', *Chem. Eur. J.* **2014**, *20*, 13099–13102.
- [2] Y. Zhu, M. Bauer, L. Ackermann, 'Late-Stage Peptide Diversification by Bioorthogonal Catalytic C–H Arylation at 23 °C in H<sub>2</sub>O', *Chem. Eur. J.* **2015**, *21*, 9980–9983.
- [3] P. S. Fier, J. F. Hartwig, 'Synthesis and Late-Stage Functionalization of Complex Molecules through C–H Fluorination and Nucleophilic Aromatic Substitution', *J. Am. Chem. Soc.* **2014**, *136*, 10139–10147.
- [4] T. Oshima, I. Yamanaka, A. Kumar, J. Yamaguchi, T. Nishiwaki-Ohkawa, K. Muto, R. Kawamura, T. Hirota, K. Yagita, S. Irle, S. A. Kay, T. Yoshimura, K. Itami, 'C–H Activation Generates Period-Shortening Molecules That Target Cryptochrome in the Mammalian Circadian Clock', *Angew. Chem. Int. Ed.* **2015**, *54*, 7193–7197.
- [5] W. R. Gutekunst, P. S. Baran, 'Total Synthesis and Structural Revision of the Piperarborenines via Sequential Cyclobutane C–H Arylation', *J. Am. Chem. Soc.* **2011**, *133*, 19076–19079.
- [6] J. Xie, C. Zhu, 'Sustainable C(sp<sup>3</sup>)-H Bond Functionalization', Springer, Heidelberg, 2016.
- [7] K. Yoshida, S. Horiuchi, N. Iwade, F. Kawagoe, T. Imamoto, 'An Efficient Route to Benzene and Phenol Derivatives via Ring-Closing Olefin Metathesis', *Synlett* **2007**, 1561–1564.
- [8] X. Yang, A. Althammer, P. Knochel, 'Selective Functionalization in Positions 2 and 3 of Indole via an Iodine-Copper Exchange Reaction', *Org. Lett.* **2004**, *6*, 1665–1667.
- [9] K.-S. Huang, E.-C. Wang, 'A novel synthesis of substituted naphthalenes via Claisen rearrangement and RCM reaction', *Tetrahedron Lett.* **2001**, *42*, 6155–6157.
- [10] K. Huang, E. Wang, 'Synthesis of Substituted Indenes from Isovanillin via Claisen Rearrangement and Ring-Closing Metathesis', *J. Chin. Chem. Soc.* **2004**, *51*, 383–391.
- [11] T. J. Donohoe, A. J. Orr, K. Gosby, M. Bingham, 'A Metathesis Approach to Aromatic Heterocycles', *Eur. J. Org. Chem.* **2005**, 1969–1971.
- [12] M.-L. Bennasar, E. Zulaica, S. Tummers, 'A synthetic entry to 2,3-fused ring indole derivatives by ring-closing metathesis reactions', *Tetrahedron Lett.* **2004**, *45*, 6283–6285.
- [13] Y. Chen, H. V. R. Dias, C. J. Lovely, 'Synthesis of fused bicyclic imidazoles by ring-closing metathesis', *Tetrahedron Lett.* **2003**, *44*, 1379–1382.
- [14] W. A. L. van Otterlo, E. L. Ngidi, E. M. Coyanis, C. B. de Koning, 'Ring-closing metathesis for the synthesis of benzo-fused bicyclic compounds', *Tetrahedron Lett.* **2003**, *44*, 311–313.
- [15] S. Kotha, K. Mandal, 'A Retrospective on the Design and Synthesis of Novel Molecules through a Strategic Consideration of Metathesis and Suzuki–Miyaura Cross-Coupling', *Chem. Asian J.* **2009**, *4*, 354–362.
- [16] S. Kotha, S. Halder, E. Brahmachary, 'Synthesis of highly functionalized phenylalanine derivatives via cross-ene metathesis reactions', *Tetrahedron* **2002**, *58*, 9203–9208.
- [17] S. Kotha, S. Halder, E. Brahmachary, T. Ganesh, 'Synthesis of Unusual  $\alpha$ -Amino Acid Derivatives via Cross-Enyne Metathesis Reaction', *Synlett* **2000**, 853–855.
- [18] S. Kotha, V. R. Shah, K. Mandal, 'Formation of Arenes via Diallylarenes: Strategic Utilization of Suzuki–Miyaura Cross-Coupling, Claisen Rearrangement and Ring-Closing Metathesis', *Adv. Synth. Catal.* **2007**, *349*, 1159–1172.
- [19] K. Yoshida, T. Toyoshima, I. Tsunoe, 'Synthesis of Aromatic Compounds Using Combinations of Ring-Closing Olefin Metathesis, Dehydration, Oxidation, and Tautomerization', *Bull. Chem. Soc. Jpn.* **2008**, *81*, 1512–1517.
- [20] S. K. Chattopadhyay, B. K. Pal, S. Maity, 'Combined Multiple Claisen Rearrangement and Ring-closing Metathesis as a Route to Naphthalene, Anthracene, and Anthracene Ring Systems', *Chem. Lett.* **2003**, *32*, 1190–1191.
- [21] V. Sabatino, J. G. Rebele, T. R. Ward, '"Close-to-Release": Spontaneous Bioorthogonal Uncaging Resulting from Ring-Closing Metathesis', *J. Am. Chem. Soc.* **2019**, *141*, 17048–17052.
- [22] B. Ferrié, S. Reymond, P. Capdevielle, J. Cossy, 'Concise Total Synthesis of (–)-Muricatacin and (–)-iso-Cladospolide B Using Chemoselective Cross-Metathesis', *Synlett* **2007**, 2891–2893.
- [23] K. Kawai, Y. Bunno, T. Yoshino, S. Matsunaga, 'Weinreb Amide Directed Versatile C–H Bond Functionalization under (η<sup>5</sup>-Pentamethylcyclopentadienyl)cobalt(III) Catalysis', *Chem. Eur. J.* **2018**, *24*, 10231–10237.
- [24] F. Li, C. Yu, J. Zhang, G. Zhong, 'Weinreb amide directed cross-coupling reaction between electron-deficient alkenes catalyzed by a rhodium catalyst', *Org. Biomol. Chem.* **2017**, *15*, 1236–1244.

Received February 10, 2021

Accepted March 30, 2021